



# Forced waves and their asymptotics in a Lotka–Volterra cooperative model under climate change<sup>☆</sup>

Yong Yang<sup>a,b</sup>, Chufen Wu<sup>b,\*</sup>, Zunxian Li<sup>c</sup>

<sup>a</sup>School of Electronics and Information Technology, Sun Yat-Sen University, Guangzhou 510006, China

<sup>b</sup>School of Mathematics and Big Data, Foshan University, Foshan 528000, China

<sup>c</sup>Department of Mathematics, Tianjin University of Technology, Tianjin 300384, China

## ARTICLE INFO

### MSC:

35K57

35C07

92D40

92D25

### Keywords:

Lotka–Volterra cooperative system

Climate change

Heterogeneity

Forced wave front

## ABSTRACT

The existence and asymptotics of forced traveling wave front in a Lotka–Volterra cooperative system under climate change are concerned in this paper. By constructing appropriate upper and lower solutions combined with the monotone iteration scheme, we show that for any given positive speed of the shifting habitat edge, there exists a nondecreasing wave front with the speed consistent with the habitat shifting speed, which indicates that no matter how moderate of the habitat shifts, two species will still be driven to extinction as the mutualistic effects exist but weak. Numerical simulations are also given to illustrate our results.

© 2019 Elsevier Inc. All rights reserved.

## 1. Introduction

Climate change such as global warming is believed to be the greatest threat to biodiversity [31]. Those species that fail to respond effectively to warming or habitat loss have been undergoing a high risk of extinction [39]. A recent report [40] illustrated that if there was a 4.5°C global mean temperature increase, then 69% of plant species would forfeit in Amazon, 89% of amphibians would turn into locally extinct in south-west Australia, up to 90% of amphibians, 86% of birds and 80% of mammals would potentially die out in the Miombo Woodlands and Southern Africa. In the wake of global warming, species are moving from their natural habitats and intend to migrate towards the poles and up elevation where is cooler. For example, umbrella trees are creeping northwards through the United States [29], tropical birds in New Guinean mountains are migrating upslope rapidly [8] and southern Africa's typical quiver tree is escaping from the equator [26]. Climate change drives the shifts in species range and distribution [4,25].

One critical question is whether a given species is capable of moving along with the continuously changing climate. For birds and certain mammals, there is at least an opportunity that they can keep pace with their ideal habitat as it shifts toward the poles or into higher altitudes. For reptiles and plants, it becomes improbable. And for those species already living in the poles-like places such as Arctic, there may be nowhere further north to go.

To characterize the heterogeneity of the shifting habitat, one effective way is to consider the environment “on the move” induced by the global climate change. Since the boundary between suitable and unsuitable habitat varies, we can take into account the favourable habitats move in the positive direction at a fixed speed of  $c$  units per timestep. In order to investigate

<sup>☆</sup> This work was supported by NSF of China [Grant nos. 11401096, 61872396 and 11601384].

\* Corresponding author.

E-mail address: [chufenwu@126.com](mailto:chufenwu@126.com) (C. Wu).

the patch moving on the persistence of a single species, Berestycki et al. [1] incorporated the climate shifts as well as the phenotypic characteristics of the population, such as dispersal ability, intrinsic growth rate, mutations and intraspecific competitions into Species Distribution Models and introduced a simplified model, taking the form of a reaction-diffusion equation

$$u_t(t, x) = du_{xx}(t, x) + g(x - ct, u(t, x)), \quad t > 0, x \in \mathbb{R}. \quad (1)$$

Here  $u(t, x)$  denotes the population density at time  $t$  and location  $x$ . The function  $g$  represents the net effect of reproduction and mortality and  $d > 0$  is the diffusion rate for species. Since the climate change drives the space to become heterogeneous, some regions are more favourable than others for the species. The growth function is therefore heterogeneous but not periodic (see [43] and references therein) in space, and the term  $g(x - ct, u(t, x))$  expresses the suitability profile of this habitat. They deduced a critical size of the favorable patch for species persistence and showed a striking asymmetry in the comoving population profile by using analytical approaches.

A traveling wave front is able to describe a species that was well distributed over some certain area would gradually expand its range to the whole environment. Global warming tendency offers chances for species to expand their ranges northwards. If the species can track the climatic change, what happens to the size and form of its population profile as time evolves? How the heterogeneities induced by climate change influence the characteristics of front propagation such as front speeds, front profiles, and front locations [20,44]? To comprehend the effect of range shifts on genetic diversity, Garnier and Lewis [10] studied the existence of traveling wave fronts to the population model (1). They decomposed the traveling wave solution into neutral genetic components and then analysed the spatio-temporal dynamics of its genetic structure. More recently, Berestycki and Fang [2] studied the complete existence and multiplicity of forced waves to (1) when  $g(s, u)$  was asymptotically of KPP type as  $s \rightarrow -\infty$ . If the reaction term in (1) is the Logistic growth function, then the population model (1) can be reduced to the Fisher's equation

$$u_t(t, x) = du_{xx}(t, x) + u(t, x)[r(x - ct) - u(t, x)], \quad t > 0, x \in \mathbb{R}. \quad (2)$$

When the suitable habitat is gradually contracting in the sense that  $c > 0$  and  $r(\cdot)$  is continuous, nondecreasing and connecting some negative number to some positive number, Hu and Zou [11] obtained the existence of extinction wave for the special model (2). Fang et al. [6] studied the existence and non-existence of KPP (monotone) wave and pulse wave to the limiting equation (2) derived from an SIS epidemic model in an expanding or contracting habitat.

Species interactions can influence the range sizes of populations directly or indirectly. A significant background problem is ascertaining when and to what extent biotic interactions really matter in determining range limits. It seems most probably that certain biotic interactions, such as obligate mutualism or generalist predation, may be crucial and needed [3,19]. Mutualism is one of the material interactions among species, which is widely existing in biosphere [28,30], especially collaborative efforts in humans to promote social development [35]. For the competitive species, climate change may be disastrous. For example, climate change drives the plants that were inhabiting previously at lower elevations move to high latitudes, which means that competition generate for alpine plants in the future. But for the cooperative species, what will it happen? The outcomes may depend on the Gambling between cooperation and climate change. A recent review elaborated the correlation of spatial structure, cooperation, mobility and cyclic interaction in evolutionary game [38], see also [24].

Consider a pair of mutualists with densities denoted by  $u_1$  and  $u_2$  and both two species follow the Logistic growth rate which is "on the move" to capture the key point that the environment is both heterogeneous and directionally shifting over time with a forced rate  $c > 0$ . The cooperative behaviors are described by  $a_i u_1 u_2$  with positive constants  $a_i$  for  $i = 1, 2$ . Then we can have the following celebrated Lotka–Volterra cooperative system

$$\begin{aligned} \partial_t u_1(t, x) &= d_1 \partial_{xx} u_1(t, x) + u_1[r_1(x - ct) - u_1 + a_1 u_2], \\ \partial_t u_2(t, x) &= d_2 \partial_{xx} u_2(t, x) + u_2[r_2(x - ct) - u_2 + a_2 u_1], \end{aligned} \quad (3)$$

where  $d_1, d_2 > 0$  are the diffusion rates. Recently, Lotka–Volterra like models have been well used in metapopulation system under heterogeneous environment such as cyclical interactions with alliance-specific heterogeneous invasion rates or stochasticity in evolutionary processes, for example, see [22,23]. A similar two species cooperative model to (3) can be also found in [3] by replacing the mutualistic effects as the Holling-II response function. Note that the cooperative model proposed in [3] has the same dynamics to (3). For the sake of simplicity and since the dynamics of (3) is sufficient to capture the main interesting properties of the spatial structure of the fronts, hereafter we only consider the traveling waves to (3).

For the existence of traveling wave fronts to some kinds of cooperative systems without considering the climate change, the reader can consult for instance [12,14–18,27,41,42] and references therein. With respect to the diffusion-cooperative system, one might consider that the transition from the origin to the stable state gives rise to the formation of traveling fronts as observed in the Fisher-KPP equation [13]. Furthermore, fronts that initiate from a local perturbation and propagate into a linearly unstable state come in two types: pulled fronts and pushed fronts [5,9,32,34], which extend the previous definitions on general transition waves. Such scenarios commonly appear in monostable reaction-diffusion systems with constant coefficients. For spatial lattice, Szolnoki et al. [36,37] found rich dynamics such as continuous and discontinuous transition waves as well as cyclic dominance happened in human cooperation model. As for the spatial-temporal models with varying coefficients, for example, like the system (3), such topics can still be true?

In this paper, we will study the forced traveling wave fronts of (3) connecting the trivial equilibrium and positive equilibrium, including the existence of fronts and their asymptotics in two tails. Such analyses may provide us some insight on the

stability studies and the category of wave fronts. The results imply that for any given positive speed of the shifting habitat edge, there exists a wave front with the speed consistent with the habitat shifting speed and the wave front decays exponentially in two tails. The wave speed to ensure the existence of front is not determined by the linearized problem around the unstable equilibrium (0,0), which may suggest that it is not linearly determined [9,33]. Such characteristics is novel in cooperative system since this type of monotone system with constant coefficients always satisfies the linear determinacy [41]. We assume

- (A1)  $r_i(x) : \mathbb{R} \rightarrow \mathbb{R}$  is continuous, nondecreasing with  $-\infty < L_i = r_i(-\infty) < 0 < r_i(+\infty) = K_i < +\infty$ ,  $i = 1, 2$ ;  
 (A2)  $a_1 a_2 < 1$ ,  $L_1(1 - a_1 a_2) + a_1(K_2 + a_2 K_1) < 0$  and  $L_2(1 - a_1 a_2) + a_2(K_1 + a_1 K_2) < 0$ ;  
 (A3)  $r_i(x)$  is continuously differentiable in  $\mathbb{R}$  and both  $r'_i(\pm\infty)$  exist,  $i = 1, 2$ .

Note that the reaction system of (3) admits four equilibria  $E_0(0, 0)$ ,  $E_1(K_1, 0)$ ,  $E_2(0, K_2)$  and  $E_*(u_1^*, u_2^*)$ , where

$$u_1^* = \frac{K_1 + a_1 K_2}{1 - a_1 a_2} > K_1, \quad u_2^* = \frac{K_2 + a_2 K_1}{1 - a_1 a_2} > K_2.$$

The rest of this article is organized as follows. In Section 2, by constructing appropriate upper and lower solutions to the coupled equations (8) and combined with the classic monotone iteration approaches developed in Wu and Zou [42], we can obtain the existence of traveling wave fronts. In Section 3, we study the asymptotical behavior of traveling wave fronts in two tails. The numerical simulation is given to illustrate our results in the last section.

## 2. Existence of traveling wave fronts

In this section, we always assume that (A1) and (A2) hold. For simplicity, denote

$$\begin{aligned} f_1(x, u_1, u_2) &= u_1[r_1(x) - u_1 + a_1 u_2], \\ f_2(x, u_1, u_2) &= u_2[r_2(x) - u_2 + a_2 u_1]. \end{aligned} \quad (4)$$

For any  $0 \leq u_1, v_1 \leq u_1^*$ ,  $0 \leq u_2, v_2 \leq u_2^*$  and  $x \in \mathbb{R}$ , we have

$$\begin{aligned} |f_1(x, u_1, u_2) - f_1(x, v_1, v_2)| &\leq \rho_1[|u_1 - v_1| + |u_2 - v_2|], \\ |f_2(x, u_1, u_2) - f_2(x, v_1, v_2)| &\leq \rho_2[|u_1 - v_1| + |u_2 - v_2|], \end{aligned} \quad (5)$$

where  $\rho_i = 2u_i^* - L_i + a_i(u_1^* + u_2^*)$ , which imply that  $f_1(x, u_1, u_2)$ ,  $f_2(x, u_1, u_2)$  are Lipschitz continuous in  $(u_1, u_2) \in [0, u_1^*] \times [0, u_2^*]$  for any  $x \in \mathbb{R}$ . Define

$$F_i(x, u_1, u_2) = \rho_i u_i + f_i(x, u_1, u_2). \quad (6)$$

Then  $F_i(x, u_1, u_2)$  is nondecreasing in  $u_i \in [0, u_i^*]$  for  $i = 1, 2$ . Let  $\mathcal{C} = C(\mathbb{R})$  be all continuous functions from  $\mathbb{R}$  to  $\mathbb{R}$  and  $\mathcal{BC} = \mathcal{C} \cap \mathcal{L}^\infty(\mathbb{R})$  be all continuous and bounded functions from  $\mathbb{R}$  to  $\mathbb{R}$ . Denote  $\mathcal{Y} = \mathcal{BC} \times \mathcal{BC}$ . For  $u = (u_1, u_2) \in \mathcal{Y}$ , employ the norm  $\|u\|_{\mathcal{Y}} = |u_1|_{\mathcal{BC}} + |u_2|_{\mathcal{BC}}$  with  $|u_i|_{\mathcal{BC}} = \sup_{x \in \mathbb{R}} |u_i(x)|$ . Then it follows from [21, Theorem 2.1] that the following conclusion holds.

**Theorem 2.1.** Consider the Cauchy problem

$$\begin{aligned} \partial_t u_1(t, x) &= d_1 \partial_{xx} u_1(t, x) - \rho_1 u_1 + F_1(x - ct, u_1, u_2), \quad t > 0, x \in \mathbb{R}, \\ \partial_t u_2(t, x) &= d_2 \partial_{xx} u_2(t, x) - \rho_2 u_2 + F_2(x - ct, u_1, u_2), \quad t > 0, x \in \mathbb{R}, \\ u(0, x) &:= (u_1(0, x), u_2(0, x)) = (u_{10}(x), u_{20}(x)) =: u_0(x), \quad x \in \mathbb{R}, \end{aligned} \quad (7)$$

where  $F_i$  is defined in (6). If  $u_0(x) \in \mathcal{Y}$  with  $0 \leq u_{i0}(x) \leq u_i^*$  in  $\mathbb{R}$ , then (7) has a unique classic solution  $u(t, \cdot) \in \mathcal{Y}$  with  $0 \leq u_i(t, x) \leq u_i^*$  for all  $t > 0$  and  $x \in \mathbb{R}$ .

Now putting  $u_i(t, x) = U_i(\xi)$  with  $\xi = x - ct$  and plugging it into (3), we have

$$\begin{aligned} d_1 U_1''(\xi) + c U_1'(\xi) + U_1(\xi)[r_1(\xi) - U_1(\xi) + a_1 U_2(\xi)] &= 0, \\ d_2 U_2''(\xi) + c U_2'(\xi) + U_2(\xi)[r_2(\xi) - U_2(\xi) + a_2 U_1(\xi)] &= 0. \end{aligned} \quad (8)$$

The solution of (8) will be considered and which is subjected to the following asymptotic boundary conditions:

$$\lim_{\xi \rightarrow -\infty} (U_1(\xi), U_2(\xi)) = (0, 0), \quad \lim_{\xi \rightarrow +\infty} (U_1(\xi), U_2(\xi)) = (u_1^*, u_2^*). \quad (9)$$

To construct the proper lower solution of (8), we introduce the following lemma, which can be found in [7]. For vector functions in  $\mathbb{R}^n$ , we will employ the norm  $\|f\| = \sum_{i=1}^n |f_i|$ . For the  $n \times n$  matrix  $A$  with elements  $a_{ij}$ , we use the norm  $\|A\| = \sum_{i,j=1}^n |a_{ij}|$ .

**Lemma 2.1.** (Poincaré–Lyapunov). Consider the equation in  $\mathbb{R}^n$

$$\dot{x} = Ax + B(t)x + P(t, x), \quad x(t_0) = x_0, \quad t \in \mathbb{R}.$$

$A$  is a constant  $n \times n$  matrix with eigenvalues which have all negative real part;  $B(t)$  is a continuous  $n \times n$  matrix with the property  $\lim_{t \rightarrow +\infty} \|B(t)\| = 0$ . The vector function  $P(t, x)$  is continuous in  $t$  and  $x$  and Lipschitz continuous in  $x$  in a neighbourhood of  $x = 0$ ; moreover we have

$$\lim_{\|x\| \rightarrow 0} \frac{\|P(t, x)\|}{\|x\|} = 0 \quad \text{uniformly in } t.$$

Then there exist constants  $\delta, \gamma > 0$  and  $M \geq 1$  such that  $\|x_0\| \leq \delta$  implies

$$\|x(t)\| \leq M\|x_0\|e^{-\gamma(t-t_0)}, \quad t \geq t_0.$$

Let  $x = (x_1, x_2, x_3, x_4)^T$  with  $x_1 = U_1, x_2 = U'_1, x_3 = U_2$  and  $x_4 = U'_2$ . Then (8) can be rewritten as the following first-order differential equations

$$\begin{aligned} x'_1(\xi) &= x_2, \\ x'_2(\xi) &= -\frac{c}{d_1}x_2 - \frac{1}{d_1}r_1(\xi)x_1 + \frac{1}{d_1}x_1^2 - \frac{a_1}{d_1}x_1x_3, \\ x'_3(\xi) &= x_4, \\ x'_4(\xi) &= -\frac{c}{d_2}x_4 - \frac{1}{d_2}r_2(\xi)x_3 + \frac{1}{d_2}x_3^2 - \frac{a_2}{d_2}x_1x_3. \end{aligned} \quad (10)$$

The corresponding vector form of (10) with initial data  $x(\xi_0) = x_0$  reads

$$x' = Ax + B(\xi)x + P(x), \quad x(\xi_0) = x_0, \quad \xi \in \mathbb{R},$$

where

$$A = \begin{bmatrix} 0 & 1 & 0 & 0 \\ -\frac{K_1}{d_1} & -\frac{c}{d_1} & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -\frac{K_2}{d_2} & -\frac{c}{d_2} \end{bmatrix}, \quad B(\xi) = \begin{bmatrix} 0 & 0 & 0 & 0 \\ \frac{1}{d_1}[K_1 - r_1(\xi)] & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{1}{d_2}[K_2 - r_2(\xi)] & 0 \end{bmatrix},$$

and  $P(x) = (0, \frac{1}{d_1}x_1(x_1 - a_1x_3), 0, \frac{1}{d_2}x_3(x_3 - a_2x_1))^T$ .

It is easy to check that  $A, B(\cdot), P(\cdot)$  satisfy the conditions in Lemma 2.1. Thus there exist constants  $\delta, \gamma > 0$  and  $M \geq 1$  such that  $\|x_0\| \leq \delta$  implies

$$\|x(\xi; \xi_0, x_0)\| \leq M\|x_0\|e^{-\gamma(\xi-\xi_0)}, \quad \xi \geq \xi_0. \quad (11)$$

Here  $x(\xi; \xi_0, x_0) = (x_1(\xi; \xi_0, x_0), x_2(\xi; \xi_0, x_0), x_3(\xi; \xi_0, x_0), x_4(\xi; \xi_0, x_0))^T$  with  $x(\xi_0; \xi_0, x_0) = x_0$ .

**Lemma 2.2.** Let  $\eta_i$  be the positive root of  $d_i\lambda^2 + c\lambda + L_i = 0$  with  $L_i = r_i(-\infty) < 0$ . Then there exist  $\alpha_i > \eta_i$  and small  $\epsilon > 0$  such that the functions

$$\underline{U}_1(\xi) = \begin{cases} \epsilon e^{\alpha_1(\xi-\xi_0)}, & \xi < \xi_0, \\ x_1(\xi; \xi_0, x_0), & \xi \geq \xi_0, \end{cases} \quad \underline{U}_2(\xi) = \begin{cases} \epsilon e^{\alpha_2(\xi-\xi_0)}, & \xi < \xi_0, \\ x_3(\xi; \xi_0, x_0), & \xi \geq \xi_0 \end{cases}$$

with  $x_0 = \epsilon(1, \alpha_1, 1, \alpha_2)^T$  are continuously differentiable in  $\mathbb{R}$  and satisfy  $0 < \underline{U}_i(\xi) \leq u_i^*$  for all  $\xi \in \mathbb{R}$  and  $i = 1, 2$ . Moreover, there holds

$$\begin{aligned} d_1 \underline{U}_1''(\xi) + c \underline{U}_1'(\xi) + \underline{U}_1(\xi)[r_1(\xi) - \underline{U}_1(\xi) + a_1 \underline{U}_2(\xi)] &\geq 0, \\ d_2 \underline{U}_2''(\xi) + c \underline{U}_2'(\xi) + \underline{U}_2(\xi)[r_2(\xi) - \underline{U}_2(\xi) + a_2 \underline{U}_1(\xi)] &\geq 0 \end{aligned} \quad (12)$$

for any  $\xi \neq \xi_0$ .

**Proof.** Define

$$\epsilon = \min_{i=1,2} \left\{ 3d_i\eta_i^2 + c\eta_i, \frac{u_i^*}{2M(1 + \eta_1 + \eta_2)} \right\},$$

where  $M \geq 1$  is defined in (11). Let  $\alpha_i$  be the positive root of the equation  $d_i\lambda^2 + c\lambda + L_i - \epsilon = 0$  with  $i = 1, 2$ . Then it is easy to check that  $\eta_i < \alpha_i \leq 2\eta_i$ . Since  $\lim_{\xi \rightarrow \xi_0^-} \underline{U}_1'(\xi) = \epsilon\alpha_1 = x_2(\xi_0; \xi_0, x_0) = \lim_{\xi \rightarrow \xi_0^+} x_1'(\xi; \xi_0, x_0)$  and  $\lim_{\xi \rightarrow \xi_0^-} \underline{U}_2'(\xi) = \epsilon\alpha_2 = x_4(\xi_0; \xi_0, x_0) = \lim_{\xi \rightarrow \xi_0^+} x_3'(\xi; \xi_0, x_0)$ , which imply  $\underline{U}_1(\xi)$  and  $\underline{U}_2(\xi)$  are continuously differentiable in  $\mathbb{R}$ . Furthermore, it follows from (4), (5) and the uniqueness of solutions to (10) subjected to the positive initial data  $x_0$  that  $\underline{U}_i(\xi) > 0$  with  $i = 1, 2$ .

If  $\xi < \xi_0$ , then  $\underline{U}_i(\xi) = \epsilon e^{\alpha_i(\xi-\xi_0)} < \epsilon \leq u_i^*$ . Also, for  $i \neq j \in \{1, 2\}$ , we get

$$d_i \underline{U}_i''(\xi) + c \underline{U}_i'(\xi) + \underline{U}_i(\xi)[r_i(\xi) - \underline{U}_i(\xi) + a_i \underline{U}_j(\xi)]$$

$$\begin{aligned}
&\geq d_i \underline{U}_i''(\xi) + c \underline{U}_i'(\xi) + \underline{U}_i(\xi)[r_i(\xi) - \underline{U}_i(\xi)] \\
&= \epsilon e^{\alpha_i(\xi - \xi_0)} [d_i \alpha_i^2 + c \alpha_i + r_i(\xi) - \epsilon e^{\alpha_i(\xi - \xi_0)}] \\
&\geq \epsilon e^{\alpha_i(\xi - \xi_0)} [d_i \alpha_i^2 + c \alpha_i + L_i - \epsilon] = 0.
\end{aligned}$$

If  $\xi > \xi_0$ , then it follows from (11) that  $\underline{U}_1(\xi) = x_1(\xi; \xi_0, x_0) \leq M \|x_0\| \leq 2M\epsilon(1 + \eta_1 + \eta_2) \leq u_1^*$ . Similarly,  $\underline{U}_2(\xi) \leq u_2^*$ . Moreover,

$$\begin{aligned}
&d_1 \underline{U}_1''(\xi) + c \underline{U}_1'(\xi) + \underline{U}_1(\xi)[r_1(\xi) - \underline{U}_1(\xi) + a_1 \underline{U}_2(\xi)] \\
&= d_1 x_2' + c x_2 + x_1[r_1(\xi) - x_1 + a_1 x_3] = 0,
\end{aligned}$$

and

$$\begin{aligned}
&d_2 \underline{U}_2''(\xi) + c \underline{U}_2'(\xi) + \underline{U}_2(\xi)[r_2(\xi) - \underline{U}_2(\xi) + a_2 \underline{U}_1(\xi)] \\
&= d_2 x_4' + c x_4 + x_3[r_2(\xi) - x_3 + a_2 x_1] = 0.
\end{aligned}$$

□

**Lemma 2.3.** Let  $\xi_1, \xi_2 \leq \xi_0$  so small that satisfy  $r_1(\xi_1) + a_1 u_2^* < 0$  and  $r_2(\xi_2) + a_2 u_1^* < 0$ . Assume  $\beta_1$  is the positive root of  $d_1 \lambda^2 + c \lambda + r_1(\xi_1) + a_1 u_2^* = 0$  and  $\beta_2$  the positive root of  $d_2 \lambda^2 + c \lambda + r_2(\xi_2) + a_2 u_1^* = 0$ . Then the functions  $\bar{U}_i(\xi) = \min \{u_i^* e^{\beta_i(\xi - \xi_i)}, u_i^*\} \geq \underline{U}_i(\xi)$  with  $i = 1, 2$  and further satisfy

$$d_1 \bar{U}_1''(\xi) + c \bar{U}_1'(\xi) + \bar{U}_1(\xi)[r_1(\xi) - \bar{U}_1(\xi) + a_1 \bar{U}_2(\xi)] \leq 0, \quad (13)$$

for any  $\xi \neq \xi_1$ :

$$d_2 \bar{U}_2''(\xi) + c \bar{U}_2'(\xi) + \bar{U}_2(\xi)[r_2(\xi) - \bar{U}_2(\xi) + a_2 \bar{U}_1(\xi)] \leq 0 \quad (14)$$

for any  $\xi \neq \xi_2$ .

**Proof.** Note that  $r_1(\xi_1) + a_1 u_2^* < 0$  and  $r_2(\xi_2) + a_2 u_1^* < 0$  can be guaranteed since  $r_1(-\infty)(1 - a_1 a_2) + a_1(K_2 + a_2 K_1) < 0$  and  $r_2(-\infty)(1 - a_1 a_2) + a_2(K_1 + a_1 K_2) < 0$ . If  $\xi > \xi_1$ , then  $\bar{U}_1(\xi) = u_1^* \geq \underline{U}_1(\xi)$ . Also,

$$\begin{aligned}
&d_1 \bar{U}_1''(\xi) + c \bar{U}_1'(\xi) + \bar{U}_1(\xi)[r_1(\xi) - \bar{U}_1(\xi) + a_1 \bar{U}_2(\xi)] \\
&\leq u_1^*[r_1(\xi) - u_1^* + a_1 u_2^*] \\
&\leq u_1^*[K_1 - u_1^* + a_1 u_2^*] = 0.
\end{aligned}$$

If  $\xi < \xi_1 \leq \xi_0$ , then  $\bar{U}_1(\xi) = u_1^* e^{\beta_1(\xi - \xi_1)} \geq \epsilon e^{\beta_1(\xi - \xi_0)} \geq \epsilon e^{\alpha_1(\xi - \xi_0)} = \underline{U}_1(\xi)$  due to  $\beta_1 < \alpha_1$ . Also,

$$\begin{aligned}
&d_1 \bar{U}_1''(\xi) + c \bar{U}_1'(\xi) + \bar{U}_1(\xi)[r_1(\xi) - \bar{U}_1(\xi) + a_1 \bar{U}_2(\xi)] \\
&\leq u_1^* e^{\beta_1(\xi - \xi_1)} [d_1 \beta_1^2 + c \beta_1 + r_1(\xi) - u_1^* e^{\beta_1(\xi - \xi_1)} + a_1 u_2^*] \\
&\leq u_1^* e^{\beta_1(\xi - \xi_1)} [d_1 \beta_1^2 + c \beta_1 + r_1(\xi_1) + a_1 u_2^*] = 0.
\end{aligned}$$

Similarly, we can show that  $\bar{U}_2(\xi) \geq \underline{U}_2(\xi)$  and (14) holds for any  $\xi \neq \xi_2$ . □

Define

$$\Gamma = \{(U_1, U_2) \in \mathcal{Y} : (\underline{U}_1, \underline{U}_2) \leq (U_1, U_2) \leq (\bar{U}_1, \bar{U}_2) \text{ in } \mathbb{R}\},$$

and

$$\begin{aligned}
F_1(U_1, U_2)(\xi) &= \rho_1 U_1(\xi) + U_1(\xi)[r_1(\xi) - U_1(\xi) + a_1 U_2(\xi)], \\
F_2(U_1, U_2)(\xi) &= \rho_2 U_2(\xi) + U_2(\xi)[r_2(\xi) - U_2(\xi) + a_2 U_1(\xi)].
\end{aligned}$$

Then  $F = (F_1, F_2)$  is nondecreasing in  $U = (U_1, U_2) \in \Gamma$ . Rewrite the system (8) as

$$\begin{aligned}
&d_1 U_1''(\xi) + c U_1'(\xi) - \rho_1 U_1(\xi) + F_1(U_1, U_2)(\xi) = 0, \\
&d_2 U_2''(\xi) + c U_2'(\xi) - \rho_2 U_2(\xi) + F_2(U_1, U_2)(\xi) = 0.
\end{aligned} \quad (15)$$

Note that  $U$  is a bounded entire solution of (15) if and only if  $U$  is continuous and satisfies the integral equation  $U = Q(U)$ , where  $Q = (Q_1, Q_2)$ ,

$$Q_i(U)(\xi) = \frac{1}{d_i(\lambda_{i2} - \lambda_{i1})} \int_{-\infty}^{+\infty} J_i(\xi - s) F_i(U)(s) ds,$$

and

$$J_i(\xi) = \begin{cases} e^{\lambda_{i2}\xi}, & \xi \leq 0, \\ e^{\lambda_{i1}\xi}, & \xi > 0, \end{cases} \quad \lambda_{ij} = \frac{-c + (-1)^j \sqrt{c^2 + 4d_i \rho_i}}{2d_i}, \quad i, j = 1, 2. \quad (16)$$

**Lemma 2.4.** The operator  $Q$  is nondecreasing and maps  $\Gamma$  into  $\Gamma$ . Moreover, if  $U \in \Gamma$  is nondecreasing in  $\xi$ , then  $Q(U)(\xi)$  is also nondecreasing in  $\xi$ .

**Proof.** For any  $V, W \in \Gamma$  with  $V \geq W$ , recalling  $0 \leq V_i, W_i \leq u_i^*$ , we obtain

$$\begin{aligned} F_1(V_1, V_2)(\xi) - F_1(W_1, W_2)(\xi) \\ = [\rho_1 + r_1(\xi) - V_1 - W_1 + a_1 V_2][V_1 - W_1] + a_1 W_1[V_2 - W_2] \geq 0. \end{aligned}$$

In a similar way,  $F_2$  is nondecreasing in  $U$ . Hence, by the nonnegativity of  $J_i$  defined in (16), we see that

$$\begin{aligned} Q_i(V)(\xi) - Q_i(W)(\xi) \\ = \frac{1}{d_i(\lambda_{i2} - \lambda_{i1})} \int_{-\infty}^{+\infty} J_i(\xi - s) [F_i(V)(s) - F_i(W)(s)] ds \geq 0. \end{aligned}$$

The above inequality implies that

$$Q(\underline{U})(\xi) \leq Q(U)(\xi) \leq Q(\bar{U})(\xi) \quad (17)$$

for all  $U \in \Gamma$  and  $\xi \in \mathbb{R}$ . We now show that  $Q$  maps  $\Gamma$  into  $\Gamma$ . For  $\xi \neq \xi_0$ , without loss of generality, we assume that  $\xi < \xi_0$ , then by (12), we have

$$\begin{aligned} Q_i(\underline{U})(\xi) &= \frac{1}{d_i(\lambda_{i2} - \lambda_{i1})} \left[ \int_{-\infty}^{\xi} e^{\lambda_{i1}(\xi-s)} F_i(\underline{U})(s) ds + \int_{\xi}^{+\infty} e^{\lambda_{i2}(\xi-s)} F_i(\underline{U})(s) ds \right] \\ &\geq \frac{1}{d_i(\lambda_{i2} - \lambda_{i1})} \left( \int_{-\infty}^{\xi} e^{\lambda_{i1}(\xi-s)} + \int_{\xi}^{\xi_0} e^{\lambda_{i2}(\xi-s)} + \int_{\xi_0}^{+\infty} e^{\lambda_{i2}(\xi-s)} \right) \\ &\quad \times [-d_i \underline{U}_i''(s) - c \underline{U}_i'(s) + \rho_i \underline{U}_i(s)] ds \\ &= \frac{1}{d_i(\lambda_{i2} - \lambda_{i1})} \left\{ d_i(\lambda_{i2} - \lambda_{i1}) \underline{U}_i(\xi) + d_i e^{\lambda_{i2}(\xi-\xi_0)} [\underline{U}_i'(\xi_0 + 0) - \underline{U}_i'(\xi_0 - 0)] \right. \\ &\quad \left. + (d_i \lambda_{i2} + c) e^{\lambda_{i2}(\xi-\xi_0)} [\underline{U}_i(\xi_0 + 0) - \underline{U}_i(\xi_0 - 0)] \right\}. \end{aligned}$$

For  $\xi > \xi_0$ , we get

$$\begin{aligned} Q_i(\underline{U})(\xi) &\geq \frac{1}{d_i(\lambda_{i2} - \lambda_{i1})} \left( \int_{-\infty}^{\xi_0} e^{\lambda_{i1}(\xi-s)} + \int_{\xi_0}^{\xi} e^{\lambda_{i1}(\xi-s)} + \int_{\xi}^{+\infty} e^{\lambda_{i2}(\xi-s)} \right) \\ &\quad \times [-d_i \underline{U}_i''(s) - c \underline{U}_i'(s) + \rho_i \underline{U}_i(s)] ds \\ &= \frac{1}{d_i(\lambda_{i2} - \lambda_{i1})} \left\{ d_i(\lambda_{i2} - \lambda_{i1}) \underline{U}_i(\xi) + d_i e^{\lambda_{i1}(\xi-\xi_0)} [\underline{U}_i'(\xi_0 + 0) - \underline{U}_i'(\xi_0 - 0)] \right. \\ &\quad \left. + (d_i \lambda_{i1} + c) e^{\lambda_{i1}(\xi-\xi_0)} [\underline{U}_i(\xi_0 + 0) - \underline{U}_i(\xi_0 - 0)] \right\}. \end{aligned}$$

Since  $\underline{U}_i(\xi)$  is continuously differentiable in  $\xi_0$ , we immediately have  $Q_i(\underline{U})(\xi) \geq \underline{U}_i(\xi)$  for  $\xi \neq \xi_0$ . By the continuity, it follows that  $Q_i(\underline{U})(\xi) \geq \underline{U}_i(\xi)$  for all  $\xi$ .

In a similar way, by using (13) and (14) in Lemma 2.3 and noticing  $\bar{U}_i(\xi_i + 0) = \bar{U}_i(\xi_i - 0)$ ,  $\bar{U}_i'(\xi_i + 0) \leq \bar{U}_i'(\xi_i - 0)$ , we can also prove that  $Q_i(\bar{U})(\xi) \leq \bar{U}_i(\xi)$  for  $\xi \neq \xi_i$ . Therefore, the continuity of  $Q$  implies  $Q_i(\bar{U})(\xi) \leq \bar{U}_i(\xi)$  for all  $\xi$ . These, together with (17), we see that  $Q$  maps  $\Gamma$  into  $\Gamma$ .

If  $U \in \Gamma$  is nondecreasing in  $\xi$ , then for any  $\xi \in \mathbb{R}$  and  $\theta > 0$ , we get

$$\begin{aligned} F_i(U)(\xi + \theta) - F_i(U)(\xi) \\ = [\rho_i + r_i(\xi) - U_i(\xi + \theta) - U_i(\xi) + a_i U_j(\xi + \theta)] [U_i(\xi + \theta) - U_i(\xi)] \\ + [r_i(\xi + \theta) - r_i(\xi)] U_i(\xi + \theta) + a_i U_i(\xi) [U_j(\xi + \theta) - U_j(\xi)] \geq 0 \end{aligned}$$

for  $i \neq j \in \{1, 2\}$ . Hence,

$$\begin{aligned} Q_i(U)(\xi + \theta) &= \frac{1}{d_i(\lambda_{i2} - \lambda_{i1})} \int_{-\infty}^{+\infty} J_i(\xi + \theta - s) F_i(U)(s) ds \\ &= \frac{1}{d_i(\lambda_{i2} - \lambda_{i1})} \int_{-\infty}^{+\infty} J_i(\xi - s) F_i(U)(s + \theta) ds \\ &\geq \frac{1}{d_i(\lambda_{i2} - \lambda_{i1})} \int_{-\infty}^{+\infty} J_i(\xi - s) F_i(U)(s) ds = Q_i(U)(\xi). \end{aligned}$$

□

**Theorem 2.2.** Assume (A1) and (A2) hold. Then (3) has a nondecreasing traveling wave front  $(u_1(t, x), u_2(t, x)) = (U_1(\xi), U_2(\xi))$  with  $\xi = x - ct$ .

**Proof.** Define the following iterations:

$$U_i^{(1)} = Q_i(\bar{U}_1, \bar{U}_2), \quad U_i^{(k+1)} = Q_i(U_1^{(k)}, U_2^{(k)}), \quad k \geq 1.$$

Since  $(\bar{U}_1, \bar{U}_2) \in \Gamma$  is nondecreasing in  $\mathbb{R}$ , Lemma 2.4 implies  $(U_1^{(k)}, U_2^{(k)}) \in \Gamma$ , and  $(U_1^{(k)}, U_2^{(k)})$  is nondecreasing in  $\xi$  for each fixed  $k = 1, 2, \dots$ , and further satisfies

$$(\underline{U}_1(\xi), \underline{U}_2(\xi)) \leq (U_1^{(k+1)}(\xi), U_2^{(k+1)}(\xi)) \leq (U_1^{(k)}(\xi), U_2^{(k)}(\xi)) \leq (\bar{U}_1(\xi), \bar{U}_2(\xi))$$

for all  $k = 1, 2, \dots$  and  $\xi \in \mathbb{R}$ . Thus  $\lim_{k \rightarrow \infty} U_i^{(k)}(\xi) = U_i(\xi)$  both exist for  $i = 1, 2$ . Obviously,  $U_i(\xi)$  is nondecreasing and nonnegative in  $\xi \in \mathbb{R}$ . Moreover, we have the following properties

$$(\underline{U}_1(\xi), \underline{U}_2(\xi)) \leq (U_1(\xi), U_2(\xi)) \leq (\bar{U}_1(\xi), \bar{U}_2(\xi)), \quad \forall \xi \in \mathbb{R}. \quad (18)$$

By the continuity of  $F_i$ , we know that  $F_i(U_1^{(k)}, U_2^{(k)})$  respectively converges point-wise to  $F_i(U_1, U_2)$ . We now show that  $(U_1, U_2)$  is a fixed-point of  $F = (F_1, F_2)$ . Since

$$|F_i(U_1^{(k)}, U_2^{(k)})| \leq \rho_i u_i^*,$$

then the Lebesgue's dominated convergence theorem yields that

$$\begin{aligned} U_i(\xi) &= \lim_{k \rightarrow \infty} U_i^{(k+1)}(\xi) \\ &= \lim_{k \rightarrow \infty} Q_i(U_1^{(k)}, U_2^{(k)})(\xi) \\ &= \lim_{k \rightarrow \infty} \frac{1}{d_i(\lambda_{i2} - \lambda_{i1})} \int_{-\infty}^{+\infty} J_i(\xi - s) F_i(U_1^{(k)}, U_2^{(k)})(s) ds \\ &= \frac{1}{d_i(\lambda_{i2} - \lambda_{i1})} \int_{-\infty}^{+\infty} J_i(\xi - s) F_i(U_1, U_2)(s) ds = Q_i(U_1, U_2)(\xi). \end{aligned}$$

Thus  $(U_1, U_2)$  is a solution of (15). Next, we prove  $(U_1, U_2)$  satisfies the asymptotical boundary conditions (9). Since

$$(\underline{U}_1(-\infty), \underline{U}_2(-\infty)) = (0, 0) = (\bar{U}_1(-\infty), \bar{U}_2(-\infty)),$$

hence by (18), we see that  $(U_1(-\infty), U_2(-\infty)) = (0, 0)$ . Note that  $U_i(\xi)$  is nondecreasing in  $\xi \in \mathbb{R}$  and  $U_i \in [0, u_i^*]$ . Therefore,  $\lim_{\xi \rightarrow +\infty} U_i(\xi)$  exists, denote it by  $\kappa_i$ , respectively. Also,  $\kappa_i \in [0, u_i^*]$  and  $\lim_{\xi \rightarrow +\infty} F_i(U_1, U_2)(\xi) = \rho_i \kappa_i + \kappa_i(K_i - \kappa_i + a_i \kappa_j)$  for  $i \neq j \in \{1, 2\}$ . In view of L'Hôpital's rule, we obtain

$$\begin{aligned} \kappa_i &= \lim_{\xi \rightarrow +\infty} U_i(\xi) = \lim_{\xi \rightarrow +\infty} Q_i(U_1, U_2)(\xi) \\ &= \lim_{\xi \rightarrow +\infty} \frac{1}{d_i(\lambda_{i2} - \lambda_{i1})} \left[ \int_{-\infty}^{\xi} e^{\lambda_{i1}(\xi-s)} F_i(U_1, U_2)(s) ds + \int_{\xi}^{+\infty} e^{\lambda_{i2}(\xi-s)} F_i(U_1, U_2)(s) ds \right] \\ &= \frac{1}{d_i(\lambda_{i2} - \lambda_{i1})} \left[ \frac{\rho_i \kappa_i + \kappa_i(K_i - \kappa_i + a_i \kappa_j)}{-\lambda_{i1}} + \frac{\rho_i \kappa_i + \kappa_i(K_i - \kappa_i + a_i \kappa_j)}{\lambda_{i2}} \right] \\ &= \frac{1}{\rho_i} [\rho_i \kappa_i + \kappa_i(K_i - \kappa_i + a_i \kappa_j)] = \kappa_i + \frac{\kappa_i(K_i - \kappa_i + a_i \kappa_j)}{\rho_i}. \end{aligned}$$

Thus,  $\kappa_i(K_i - \kappa_i + a_i \kappa_j) = 0$  for  $i \neq j \in \{1, 2\}$ . Since  $\kappa_i \in [0, u_i^*]$ , it must be  $\kappa_i = 0$  or  $K_i - \kappa_i + a_i \kappa_j = 0$ . If  $\kappa_i = 0$ , then since  $U_i$  is nondecreasing in  $\mathbb{R}$  and  $U_i(\pm\infty) = 0$ , we know that  $U_i \equiv 0$  in  $\mathbb{R}$ . This contradicts  $U_i(\xi) \geq \underline{U}_i(\xi)$  and the definition of  $\underline{U}_i$ . Thus,  $K_i - \kappa_i + a_i \kappa_j = 0$ , which is equivalent to  $K_1 - \kappa_1 + a_1 \kappa_2 = 0$  and  $K_2 - \kappa_2 + a_2 \kappa_1 = 0$ . Some trivial calculations show that  $\kappa_1 = u_1^*$  and  $\kappa_2 = u_2^*$ . The asymptotic boundary conditions (9) are satisfied.  $\square$

### 3. Asymptotic behaviors of traveling waves

To obtain more asymptotic information of the traveling waves at  $\pm\infty$ , we let

$$(U_1(\xi), U_2(\xi)) = (u_1^0(\xi), u_2^0(\xi)) \quad \text{for } -\infty < \xi < +\infty,$$

which is a solution of (8) satisfies (9). Differentiating (8) with respect to  $\xi$ , then we know that  $(U_1'(\xi), U_2'(\xi)) =: (\omega_1, \omega_2)$  satisfies

$$\begin{aligned} d_1 \omega_1'(\xi) + c \omega_1'(\xi) + [r_1(\xi) - 2u_1^0 + a_1 u_2^0] \omega_1 + a_1 u_1^0 \omega_2 + r_1'(\xi) u_1^0 &= 0, \\ d_2 \omega_2'(\xi) + c \omega_2'(\xi) + [r_2(\xi) - 2u_2^0 + a_2 u_1^0] \omega_2 + a_2 u_2^0 \omega_1 + r_2'(\xi) u_2^0 &= 0. \end{aligned} \quad (19)$$

**Theorem 3.1.** Assume (A1) to (A3) hold. Then there exist positive constants  $A_i, B_i, C_i$  with  $i = 1, 2$  such that the traveling wave front  $(U_1(\xi), U_2(\xi))$  to (3) has the following asymptotic properties

$$\begin{pmatrix} U_1(\xi) \\ U_2(\xi) \end{pmatrix} = \begin{pmatrix} (A_1 + o(1))e^{-\frac{c+\sqrt{c^2-4d_1L_1}}{2d_1}\xi} \\ (A_2 + o(1))e^{-\frac{c+\sqrt{c^2-4d_2L_2}}{2d_2}\xi} \end{pmatrix}$$

as  $\xi \rightarrow -\infty$ ; and

$$\begin{pmatrix} U_1(\xi) \\ U_2(\xi) \end{pmatrix} = \begin{pmatrix} u_1^* - (B_1 + o(1))e^{\lambda_2\xi} - (C_1 + o(1))e^{\lambda_4\xi} \\ u_2^* - (B_2 + o(1))e^{\lambda_2\xi} - (C_2 + o(1))e^{\lambda_4\xi} \end{pmatrix},$$

as  $\xi \rightarrow +\infty$  provided  $d_1 = d_2$ . Here  $\lambda_2, \lambda_4$  are defined in (25).

**Proof.** The limiting equations for (19) as  $\xi \rightarrow -\infty$  read

$$\begin{aligned} d_1\phi_1''(\xi) + c\phi_1'(\xi) + L_1\phi_1(\xi) &= 0, \\ d_2\phi_2''(\xi) + c\phi_2'(\xi) + L_2\phi_2(\xi) &= 0. \end{aligned} \quad (20)$$

The first equation of (20) has two independent solutions in the following form

$$\phi_1^1(\xi) = e^{-\frac{c-\sqrt{c^2-4d_1L_1}}{2d_1}\xi}, \quad \phi_1^2(\xi) = e^{-\frac{c+\sqrt{c^2-4d_1L_1}}{2d_1}\xi}.$$

Relating (19) and (20), we see that  $\omega_1$  admits the following property as  $\xi \rightarrow -\infty$

$$\omega_1(\xi) = p_1[1 + o(1)]\phi_1^1(\xi) + q_1[1 + o(1)]\phi_1^2(\xi).$$

Since  $\lim_{\xi \rightarrow -\infty} \omega_1(\xi) = 0$ , it must be  $p_1 = 0$ . Hence, for  $\xi \rightarrow -\infty$ ,

$$U_1(\xi) = \omega_1(\xi) = q_1[1 + o(1)]e^{-\frac{c+\sqrt{c^2-4d_1L_1}}{2d_1}\xi}.$$

Similarly,

$$U_2(\xi) = \omega_2(\xi) = q_2[1 + o(1)]e^{-\frac{c+\sqrt{c^2-4d_2L_2}}{2d_2}\xi},$$

as  $\xi \rightarrow -\infty$ . By integration from  $-\infty$  to  $\xi$ , it follows that

$$\begin{pmatrix} U_1(\xi) \\ U_2(\xi) \end{pmatrix} = \begin{pmatrix} (A_1 + o(1))e^{-\frac{c+\sqrt{c^2-4d_1L_1}}{2d_1}\xi} \\ (A_2 + o(1))e^{-\frac{c+\sqrt{c^2-4d_2L_2}}{2d_2}\xi} \end{pmatrix}$$

as  $\xi \rightarrow -\infty$  for some constants  $A_1, A_2 > 0$ . Now the limiting equations for (19) as  $\xi \rightarrow +\infty$  are

$$\begin{aligned} d_1\psi_1''(\xi) + c\psi_1'(\xi) + A\psi_1(\xi) + B\psi_2(\xi) &= 0, \\ d_2\psi_2''(\xi) + c\psi_2'(\xi) + C\psi_1(\xi) + D\psi_2(\xi) &= 0, \end{aligned} \quad (21)$$

where

$$A = K_1 - 2u_1^* + a_1u_2^* = -u_1^* < 0, \quad B = a_1u_1^* > 0,$$

$$D = K_2 - 2u_2^* + a_2u_1^* = -u_2^* < 0, \quad C = a_2u_2^* > 0.$$

Note that in (21) we have used the fact that  $r_i'(+\infty) = 0$  with  $i = 1, 2$ . Indeed, by applying L'Hôpital's rule, we get

$$\lim_{\xi \rightarrow +\infty} r_i(\xi) = \lim_{\xi \rightarrow +\infty} \frac{e^\xi r_i(\xi)}{e^\xi} = \lim_{\xi \rightarrow +\infty} \frac{e^\xi [r_i(\xi) + r_i'(\xi)]}{e^\xi} = \lim_{\xi \rightarrow +\infty} [r_i(\xi) + r_i'(\xi)],$$

which implies  $r_i'(+\infty) = 0$ . Letting  $\psi_1' = \tilde{\psi}_1, \psi_2' = \tilde{\psi}_2$ , then we can rewrite (21) as the following first-order differential equations

$$\begin{aligned} \psi_1'(\xi) &= \tilde{\psi}_1, \\ \tilde{\psi}_1'(\xi) &= -\frac{c}{d_1}\tilde{\psi}_1 - \frac{A}{d_1}\psi_1 - \frac{B}{d_1}\psi_2, \\ \psi_2'(\xi) &= \tilde{\psi}_2, \\ \tilde{\psi}_2'(\xi) &= -\frac{c}{d_2}\tilde{\psi}_2 - \frac{C}{d_2}\psi_1 - \frac{D}{d_2}\psi_2. \end{aligned} \quad (22)$$



The characteristic equation of (22) is

$$\lambda^2 \left( \lambda + \frac{c}{d_1} \right) \left( \lambda + \frac{c}{d_2} \right) + \frac{D}{d_2} \lambda \left( \lambda + \frac{c}{d_1} \right) + \frac{A}{d_1} \lambda \left( \lambda + \frac{c}{d_2} \right) + \frac{AD - BC}{d_1 d_2} = 0. \quad (23)$$

If  $d_1 = d_2 = d$ , then (23) can be simplified as

$$\lambda^2 \left( \lambda + \frac{c}{d} \right)^2 + \frac{1}{d} (A + D) \lambda \left( \lambda + \frac{c}{d} \right) + \frac{1}{d^2} (AD - BC) = 0.$$

Let  $\lambda \left( \lambda + \frac{c}{d} \right) = y$ . Then  $y$  satisfies  $y^2 + \frac{1}{d} (A + D) y + \frac{1}{d^2} (AD - BC) = 0$ . Since

$$(A + D)^2 - 4(AD - BC) = (A - D)^2 + 4BC > 0,$$

$$AD = u_1^* u_2^* > a_1 a_2 u_1^* u_2^* = BC, \quad A + D < 0,$$

we have  $y_j = \frac{-(A+D) + (-1)^j \sqrt{(A+D)^2 - 4(AD-BC)}}{2d} > 0$  with  $j = 1, 2$ . Thus the general solution corresponding to (22) can be expressed as

$$(\psi_1, \tilde{\psi}_1, \psi_2, \tilde{\psi}_2)^T = \ell_1 h_1 e^{\lambda_1 \xi} + \ell_2 h_2 e^{\lambda_2 \xi} + \ell_3 h_3 e^{\lambda_3 \xi} + \ell_4 h_4 e^{\lambda_4 \xi}, \quad (24)$$

where

$$\begin{aligned} \lambda_1 &= \frac{-c + \sqrt{c^2 + 4d^2 y_1}}{2d} > 0, & \lambda_2 &= \frac{-c - \sqrt{c^2 + 4d^2 y_1}}{2d} < 0 \\ \lambda_3 &= \frac{-c + \sqrt{c^2 + 4d^2 y_2}}{2d} > 0, & \lambda_4 &= \frac{-c - \sqrt{c^2 + 4d^2 y_2}}{2d} < 0, \end{aligned} \quad (25)$$

and  $h_i$  are eigenvectors corresponding to  $\lambda_i$ ,  $\ell_i$  are arbitrary constants with  $i = 1, 2, 3, 4$ . Since  $(\psi_1, \tilde{\psi}_1, \psi_2, \tilde{\psi}_2)^T \rightarrow (0, 0, 0, 0)$  as  $\xi \rightarrow +\infty$ , we follow from (24) that  $\ell_1 = 0$  and  $\ell_3 = 0$ . Thus,  $(\psi_1, \tilde{\psi}_1, \psi_2, \tilde{\psi}_2)^T = \ell_2 h_2 e^{\lambda_2 \xi} + \ell_4 h_4 e^{\lambda_4 \xi}$ . Hence,  $(\omega_1, \omega_2)$  possesses the following property as  $\xi \rightarrow +\infty$

$$\begin{pmatrix} \omega_1(\xi) \\ \omega_2(\xi) \end{pmatrix} = \begin{pmatrix} \nu_1(m_1 + o(1))e^{\lambda_2 \xi} + \nu_2(m_2 + o(1))e^{\lambda_4 \xi} \\ \nu_1(n_1 + o(1))e^{\lambda_2 \xi} + \nu_2(n_2 + o(1))e^{\lambda_4 \xi} \end{pmatrix},$$

where  $m_i, n_i, \nu_i$  are constants and  $\nu_i$  cannot be zero simultaneously,  $i = 1, 2$ . Indeed,  $m_i \neq 0$  and  $n_i \neq 0$ . For the solution  $h_i e^{\lambda_i \xi}$  to (22), if one of the first and third components of the eigenvectors  $h_i$  is zero, then the linear system (22) leads to the other components are also zero. By integrating from  $\xi$  to  $+\infty$ , it follows that


$$\begin{pmatrix} U_1(\xi) \\ U_2(\xi) \end{pmatrix} = \begin{pmatrix} u_1^* - (B_1 + o(1))e^{\lambda_2 \xi} - (C_1 + o(1))e^{\lambda_4 \xi} \\ u_2^* - (B_2 + o(1))e^{\lambda_2 \xi} - (C_2 + o(1))e^{\lambda_4 \xi} \end{pmatrix},$$

as  $\xi \rightarrow +\infty$  and  $d_1 = d_2$ .  $\square$

#### 4. Simulations

Truncate  $\mathbb{R} = (-\infty, \infty)$  by  $[-X, X]$  for some large  $X$  and adopt the uniform partition of  $[-X, X]$  as

$$-X = \xi_1 < \xi_2 < \cdots < \xi_{2n-1} < \xi_{2n} < \xi_{2n+1} = X,$$



**Fig. 1.** Illustration of monotone forced wave fronts in the discrete system (26) corresponding to wave profile system (8) with Logistic growth functions  $r_1(x) = \frac{2}{1+e^{-x}} - 1.5$  and  $r_2(x) = \frac{2}{\pi} \arctan x$ . Parameters are  $a_1 = 0.3$ ,  $a_2 = 0.5$ ,  $d_1 = d_2 = 1$ ,  $c = 2$ ,  $h = 0.1$ . The boundary condition is the characteristic function on  $(-\infty, 0)$ . The wave front of  $U_1$ -species is shown in the left plot and  $U_2$ -species in the right plot with every 10 time steps.

where  $\xi_i = \xi_1 + (i-1)h$ ,  $h = \frac{x}{n}$ ,  $i = 1, 2, \dots, 2n+1$ . Corresponding to the truncation, then the asymptotic boundary conditions (9) become to

$$(U_1(\xi_1), U_2(\xi_1)) = (0, 0), (U_1(\xi_{2n+1}), U_2(\xi_{2n+1})) = (u_1^*, u_2^*).$$

By using the conventional numeric differentiation, we get a discrete system corresponding to (8) as follows

$$\begin{aligned} (-ch + d_1)U_1(\xi_{i-1}) - (-ch + 2d_1)U_1(\xi_i) + d_1U_1(\xi_{i+1}) + h^2U_1(\xi_i)[r_1(\xi_i) - U_1(\xi_i) + a_1U_2(\xi_i)] &= 0, \\ (-ch + d_2)U_2(\xi_{i-1}) - (-ch + 2d_2)U_2(\xi_i) + d_2U_2(\xi_{i+1}) + h^2U_2(\xi_i)[r_2(\xi_i) - U_2(\xi_i) + a_2U_1(\xi_i)] &= 0. \end{aligned} \quad (26)$$

Let  $r_1(x) = \frac{2}{1+e^{-x}} - 1.5$ ,  $r_2(x) = \frac{2}{\pi} \arctan x$ ,  $a_1 = 0.3$ ,  $a_2 = 0.5$ ,  $d_1 = d_2 = 1$ ,  $c = 2$ ,  $h = 0.1$ . Then  $(u_1^*, u_2^*) \approx (0.9412, 1.4706)$  and (A1) to (A3) hold. From Fig. 1, we see that the forced wave fronts are exponentially decaying in two tails.

## Acknowledgments

The authors would like to thank the associate editor and the referees for their very helpful comments and suggestions.

## Supplementary material

Supplementary material associated with this article can be found, in the online version, at doi:10.1016/j.amc.2019.01.058.

## References

- [1] H. Berestycki, O. Diekmann, C.J. Nagelkerke, P.A. Zegeling, Can a species keep pace with a shifting climate, *Bull. Math. Biol.* 71 (2009) 399–429.
- [2] H. Berestycki, J. Fang, Forced waves of the fisher-KPP equation in a shifting environment, *J. Differ. Equ.* 264 (2018) 2157–2183.
- [3] C. Cosner, Challenges in modeling biological invasions and population distributions in a changing climate, *Ecol. Complex.* 20 (2014) 258–263.
- [4] M.B. Davis, R.G. Shaw, Range shifts and adaptive responses to quaternary climate change, *Science* 292 (2001) 673–679.
- [5] U. Ebert, W. van Saarloos, Front propagation into unstable states: universal algebraic convergence towards uniformly translating pulled fronts, *Physica D* 146 (2000) 1–99.
- [6] J. Fang, Y.J. Lou, J. Wu, Can pathogen spread keep pace with its host invasion? *SIAM J. Appl. Math.* 76 (4) (2016) 1633–1657.
- [7] F. Verhulst, in: *Nonlinear Differential Equations and Dynamical Systems*, 2nd ed., Universitext, Springer-Verlag, Berlin, 1996.
- [8] B.G. Freeman, A.M.C. Freeman, Rapid upslope shifts in new guinean birds illustrate strong distributional responses of tropical montane species to global warming, *Proc. Natl. Acad. Sci. USA* 111 (2014) 4490–4494.
- [9] J. Garnier, T. Giletti, F. Hamel, L. Roques, Inside dynamics of pulled and pushed fronts, *J. Math. Pures Appl.* 98 (2012) 428–449.
- [10] J. Garnier, M. Lewis, Expansion under climate change: the genetic consequences, *Bull. Math. Biol.* 78 (2016) 2165–2185.
- [11] H.J. Hu, X. Zou, Existence of an extinction wave in the fisher equation with a shifting habitat, *Proc. Am. Math. Soc.* 145 (2017) 4763–4771.
- [12] J.H. Huang, X. Zou, Traveling wavefronts in diffusive and cooperative Lotka–Volterra system with delays, *J. Math. Anal. Appl.* 271 (2002) 455–466.
- [13] M. Iida, R. Lui, H. Ninomiya, Stacked fronts for cooperative systems with equal diffusion coefficients, *SIAM J. Math. Anal.* 43 (2011) 1369–1389.
- [14] W.T. Li, Z.C. Wang, Traveling fronts in diffusive and cooperative Lotka–Volterra system with nonlocal delays, *Z. Angew. Math. Phys.* 58 (2007) 571–591.
- [15] B. Li, L. Zhang, Travelling wave solutions in delayed cooperative systems, *Nonlinearity* 24 (2011) 1759–1776.
- [16] B. Li, Traveling wave solutions in partially degenerate cooperative reaction–diffusion systems, *J. Differ. Equ.* 252 (2012) 4842–4861.
- [17] X. Liang, X.Q. Zhao, Asymptotic speeds of spread and traveling waves for monotone semiflows with applications, *Commun. Pure Appl. Math.* 60 (2007) 1–40. *Comm. Pure Appl. Math.* 61 (2008) 137–138 (Erratum).
- [18] G. Lin, W.T. Li, S. Ruan, Monostable wavefronts in cooperative Lotka–Volterra systems with nonlocal delays, *Discret. Contin. Dyn. Sys. A* 31 (2011) 1–23.
- [19] X.B. Lin, P. Weng, C.F. Wu, Traveling wave solutions for a predator–prey system with sigmoidal response function, *J. Dyn. Differ. Equ.* 23 (2011) 903–921.
- [20] F. Lutscher, E. Pachepsky, M.A. Lewis, The effect of dispersal patterns on stream populations, *SIAM Rev.* 47 (2005) 749–772.
- [21] C.V. Pao, *Nonlinear Parabolic and Elliptic Equations*, Plenum Press, New York, 1992.
- [22] M. Perc, A. Szolnoki, G. Szabó, Cyclical interactions with alliance-specific heterogeneous invasion rates, *Phys. Rev. E* 75 (2007) 052102.
- [23] M. Perc, A. Szolnoki, Noise-guided evolution within cyclical interactions, *New J. Phys.* 9 (2007) 267.
- [24] M. Perc, P. Grigolini, Collective behavior and evolutionary games—an introduction, *Chaos Solitons Fractals* 56 (2013) 1–5.
- [25] A.L. Perry, P.J. Low, J.R. Ellis, J.D. Reynolds, Climate change and distribution shifts in marine fishes, *Science* 308 (2005) 1912–1915.
- [26] G.F. Midgley, W.J. Bond, Future of african terrestrial biodiversity and ecosystems under anthropogenic climate change, *Nat. Clim. Change* 5 (2015) 823–829.
- [27] K. Mischaikow, V. Hutson, Travelling waves for mutualist species, *SIAM J. Math. Anal.* 24 (1993) 987–1008.
- [28] J. Murray, *Mathematical Biology*, Springer, New York, 1989.
- [29] D.L. Murray, M.J.L. Peers, Y.N. Majchrzak, M. Wehtje, C. Ferreira, R.S.A. Pickles, J.R. Row, D.H. Thornton, Continental divide: predicting climate-mediated fragmentation and biodiversity loss in the boreal forest, *PLoS ONE* 12 (5) (2017) e0176706.
- [30] Y. Nakata, Y. Muroya, Permanence for nonautonomous Lotka–Volterra cooperative systems with delays, *Nonlinear Anal. Real World Appl.* 11 (2010) 528–534.
- [31] T. Newbold, Future effects of climate and land-use change on terrestrial vertebrate community diversity under different scenarios, *Proc. R. Soc. B* 285 (2018) 20180792.
- [32] F. Rothe, Convergence to pushed fronts, *Rocky Mt. J. Math.* 11 (1981) 617–634.
- [33] A. Solar, S. Trofimchuk, Asymptotic convergence to pushed wavefronts in a monostable equation with delayed reaction, *Nonlinearity* 28 (2015) 2027–2052.
- [34] A.N. Stokes, On two types of moving front in quasilinear diffusion, *Math. Biosci.* 31 (1976) 307–315.
- [35] A. Szolnoki, M. Perc, Reward and cooperation in the spatial public goods game, *EPL* 92 (2010) 38003.
- [36] A. Szolnoki, M. Perc, Correlation of positive and negative reciprocity fails to confer an evolutionary advantage: phase transitions to elementary strategies, *Phys. Rev. X* 3 (2013) 041021.
- [37] A. Szolnoki, M. Perc, G. Szabó, Defense mechanisms of empathetic players in the spatial ultimatum game, *Phys. Rev. Lett.* 109 (2012) 078701.
- [38] A. Szolnoki, M. Mobilia, L.L. Jiang, B. Szczesny, A.M. Rucklidge, M. Perc, Cyclic dominance in evolutionary games: a review, *J. R. Soc. Interface* 11 (2014) 20140735.
- [39] C.D. Thomas, A. Cameron, R.E. Green, et al., Extinction risk from climate change, *Nature* 427 (2004) 145–148.

- [40] R. Warren, J. Price, J. VanDerWal, S. Cornelius, H. Sohl, The implications of the united nations paris agreement on climate change for globally significant biodiversity areas, *Clim. Change* 147 (2018) 395–409.
- [41] H. Weinberger, M. Lewis, B. Li, Analysis of linear conjecture for spread in cooperative models, *J. Math. Biol.* 45 (2002) 183–218.
- [42] J. Wu, X. Zou, Traveling wave fronts of reaction-diffusion systems with delay, *J. Dyn. Differ. Equ.* 13 (2001) 651–686.
- [43] C.F. Wu, D.M. Xiao, X.Q. Zhao, Spreading speeds of a partially degenerate reaction-diffusion system in a periodic habitat, *J. Differ. Equ.* 255 (2013) 3893–4011.
- [44] J. Xin, Front propagation in heterogeneous media, *SIAM Rev.* 42 (2000) 161–230.